

McGill University
Department of Mathematics and Statistics

Ph.D. preliminary examination, PART A

PURE MATHEMATICS
Paper BETA

19 August 2016
1:00 p.m. - 5:00 p.m.

INSTRUCTIONS:

(i) This paper consists of the three modules (1) Algebra, (2) Analysis, and (3) Geometry & Topology, each of which comprises 4 questions. You should answer 7 questions with at least 2 from each module.

(ii) Pay careful attention to the exposition. Make an effort to ensure that your arguments are

Algebra Module

[ALG. 1]

Analysis Module

[AN. 1]

Let $\epsilon > 0$ be fixed. Show that the set of all real numbers $x \in [0; 1]$ such that there exist infinitely many pairs $p, q \in \mathbf{N}$ such that $|x - \frac{p}{q}| < \frac{1}{q^2 + \epsilon}$ has Lebesgue measure 0.

[AN. 2]

Let f be a uniformly continuous function on \mathbb{R} . Suppose that $f \in L^p$ for some $p, 1 < p < \infty$. Prove that $f(x) \neq 0$ as $|x| \rightarrow \infty$.

[AN. 3]

(a) Give a definition of $\|f\|_1$ of a measurable complex function f .

(b) Recall that the essential range of a function $f \in L^1(\mathbb{R}; \mathbb{C})$ is the set consisting of complex numbers w such that

$$(\{x : |f(x) - w| < \epsilon\}) > 0$$

for every $\epsilon > 0$. Prove that R_f is compact.

(c) Show that $\|f\|_1 = \sup_{w \in R_f} \int |f - w|$.

[AN. 4]

(a) Give a definition of a locally compact topological space.

(b) Give an example of a Borel measure on \mathbb{R} such that $X = L^2(\mathbb{R}; \mu)$ is locally compact and explain why it is so.

(c) Give an example of a Borel measure on \mathbb{R} such that $X = L^2(\mathbb{R}; \mu)$ is not locally compact and explain why it is so.

Geometry and Topology Module

[GT. 1]

(a) Suppose that X is a separable metric space. Show that any subspace of X is separable.

(b) Suppose that X is a compact metric space. Show that X is separable and that any compatible metric on X is complete.

[GT. 2]

(a) Show that the connected sum $T \# P$ of the torus T and the projective plane P is homeomorphic to the connected sum of three copies of the projective plane $P \# P \# P$.

(b) The boundary of the Möbius band is a circle. Which surface do we obtain if we identify antipodal points of that circle? Justify your answer.

[GT. 3]

Let G be a Lie group acting on a manifold M transitively, let H be a connected compact Lie subgroup of G which is an isotropic group of a point $p \in M$. Show that M has a Riemannian metric such that the transformation determined by each element of G is an isometry.

[GT. 4]

Let M be a Riemannian manifold of dimension n and let $p \in M$. Prove that there is a neighborhood U of p and n vector fields e_1, \dots, e_n in U , such that

$$\langle e_i, e_j \rangle = \delta_{ij}; \quad \nabla_{e_i} e_j(p) = 0; \quad \delta_{ij} = 1; \quad i, j = 1, \dots, n;$$